



ᑲᑭᑎᑭᑭᑭᑭᑭᑭ ᑭᑭᑭᑭᑭᑭᑭᑭᑭᑭᑭᑭᑭᑭ  
Building *Nunavut* Together  
*Nunavut* iuqatigiingniq  
Bâtir le *Nunavut* ensemble

# Public Service Announcement

## Syphilis is on the rise in Nunavut

**Start Date: December 11, 2013**

**End Date: January 31, 2014**

**Nunavut**

**60 sec**

---

Syphilis is a sexually transmitted infection (STI) that is on the rise in Nunavut.

Syphilis is a serious STI that is passed from one person to another during sex. Syphilis can also be passed from mother to child during birth.

An early symptom is a painless sore that usually appears three to 90 days after sexual contact with an infected person. Many people don't notice the sore because it is painless.

It is important to be tested if you think you have syphilis. Syphilis is easily treated with antibiotics if caught early. However, over time, if left untreated it can cause severe damage to the heart, blood vessels and brain, and even death. People with syphilis are also at greater risk of contracting HIV/AIDS.

The more sexual partners you have, the greater the risk that you will be exposed to an STI. If you participate in excessive drinking or drug use, you reduce your inhibitions and may participate in unprotected sex. Be careful – using a condom every time you have sex can greatly reduce the risk of getting an STI.

Please contact your health care worker if you think you have syphilis. Getting tested for an STI is easy and it's confidential.

###

